

REFERENCES

References to studies included in this review

Banks 2004 {published data only}

Banks P, Wright J, O'Brien K. Incremental versus maximum bite advancement during twin-block therapy: a randomized controlled clinical trial. *American Journal of Orthodontics and Dentofacial Orthopedics* 2004;**126**(5):583–8.

Bilgiç 2011 {published data only}

Bilgiç F, Hamamci O, Baş arın G. Comparison of the effects of fixed and removable functional appliances on the skeletal and dentoalveolar structures. *Australian Orthodontic Journal* 2011;**27**(2):110–6.

Cevidaneş 2003 {published data only}

Cevidaneş LH, Franco AA, Scanavini MA, Vigorito JW, Enlow DH, Proffit WR. Clinical outcomes of Frankel appliance therapy assessed with a counterpart analysis. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003;**123**(4):379–87.

Cura 1997 {published data only}

Cura N, Sarac M. The effect of treatment with the Bass appliance on skeletal Class II malocclusions: a cephalometric investigation. *European Journal of Orthodontics* 1997;**19**(6):691–702.

Florida 1998 {published and unpublished data}

Chen DR, McGorray SP, Dolce C, Wheeler TT. Effect of early Class II treatment on the incidence of incisor trauma. *American Journal of Orthodontics and Dentofacial Orthopedics* 2011;**140**(4):e155–60.

Dolce C, McGorray SP, Brazeau L, King GJ, Wheeler TT. Timing of Class II treatment: skeletal changes comparing 1-phase and 2-phase treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 2007; Vol. 132, issue 4:481–9.

Johnson PD, Cohen DA, Aiosa L, McGorray S, Wheeler T. Attitudes and compliance of pre-adolescent children during early treatment of Class II malocclusion. *Clinical Orthodontics and Research* 1998;**1**(1):20–8.

Keeling SD, Garvan CW, King GJ, Wheeler TT, McGorray S. Temporomandibular disorders after early Class II treatment with bionators and headgears: results from a randomized controlled trial. *Seminars in Orthodontics* 1995;**1**(3):149–64.

* Keeling SD, Wheeler TT, King GJ, Garvan CW, Cohen DA, Cabassa S, et al. Anteroposterior skeletal and dental changes after early Class II treatment with bionators and headgear. *American Journal of Orthodontics and Dentofacial Orthopedics* 1998;**113**(1):40–50.

King GJ, McGorray SP, Wheeler TT, Dolce C, Taylor M. Comparison of peer assessment ratings (PAR) from 1-phase and 2-phase treatment protocols for Class II malocclusions. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003;**123**(5):489–96.

King GJ, Wheeler TT, McGorray SP. Randomised prospective clinical trial evaluating early treatment of Class

II malocclusions. *European Journal of Orthodontics* 1999;**21**(4):445.

Wheeler TT, McGorray SP, Dolce C, Taylor MG, King GJ. Effectiveness of early treatment of Class II malocclusion. *American Journal of Orthodontics and Dentofacial Orthopedics* 2002;**121**(1):9–17.

Wortham JR, McGorray S, Taylor M, Dolce C, King DJ, Wheeler TT. Arch dimension changes following phase I and phase II orthodontic class II treatment. *Journal of Dental Research* 2001;**80**(Spec Issue (AADR Abstracts)):177 (Abs No 1131).

Ghafari 1998 {published data only}

Ghafari J. Class II malocclusion: Comparison of alternative treatments and time of treatment. *European Journal of Orthodontics* 1999;**21**(4):439–40.

Ghafari J, Efstratiadis S, Shofer FS, Markowitz D, Jacobsson-Hunt U, Baumrind S, et al. Relation between occlusal and cephalometric changes in the treatment of distocclusion. *Journal of Dental Research* 1999;**78**(March Spec Issue):443 (Abs No 2699).

Ghafari J, Jacobsson-Hunt U, Markowitz DL, Shofer FS, Laster LL. Changes of arch width in the early treatment of Class II, division 1 malocclusions. *American Journal of Orthodontics and Dentofacial Orthopedics* 1994;**106**(5):496–502.

Ghafari J, King GJ, Tulloch JF. Early treatment of Class II, division 1 malocclusion—comparison of alternative treatment modalities. *Clinical Orthodontics and Research* 1998;**1**(2):107–17.

* Ghafari J, Shofer FS, Jacobsson-Hunt U, Markowitz DL, Laster LL. Headgear versus function regulator in the early treatment of Class II, division 1 malocclusion: a randomized clinical trial. *American Journal of Orthodontics and Dentofacial Orthopedics* 1998;**113**(1):51–61.

Ghafari JG, Shofer FS, Laster LL, Markowitz DL, Silverton S, Katz SH. Monitoring growth during orthodontic treatment. *Seminars in Orthodontics* 1995;**1**(3):165–75.

Jamilian 2011 {published and unpublished data}

Jamilian A, Showkatbakhsh R, Amiri SS. Treatment effects of the R-appliance and twin block in Class II division 1 malocclusion. *European Journal of Orthodontics* 2011;**33**(4):354–8.

Lee 2007 {published data only}

Lee RT, Kyi CS, Mack GJ. A controlled clinical trial of the effects of the Twin Block and Dynamax appliances on the hard and soft tissues. *European Journal of Orthodontics* 2007;**29**(3):272–82.

London 1998 {published data only}

* Illing HM, Morris DO, Lee RT. A prospective evaluation of Bass, Bionator and Twin Block appliances. Part I—The hard tissues. *European Journal of Orthodontics* 1998;**20**(5):501–16.

McDonagh S, Moss JP, Goodwin P, Lee RT. A prospective optical surface scanning and cephalometric assessment

- of the effect of functional appliances on the soft tissues. *European Journal of Orthodontics* 2001;**23**(2):115–26.
- Morris DO, Illing HM, Lee RT. A prospective evaluation of Bass, Bionator and Twin Block appliances. Part II--The soft tissues. *European Journal of Orthodontics* 1998;**20**(6): 663–84.
- Mao 1997** {published data only}
Mao J, Zhao H. The correction of Class II, division 1 malocclusion with bionator headgear combination appliance. *Journal of Tongji Medical University* 1997;**17**(4): 254–6.
- New Zealand 2000** {published data only}
Courtney M, Harkness M, Herbison P. Maxillary and cranial base changes during treatment with functional appliances. *American Journal of Orthodontics and Dentofacial Orthopedics* 1996;**109**(6):616–24.
Nelson C, Harkness M, Herbison P. Mandibular changes during functional appliance treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 1993;**104**(2): 153–61.
O'Neill K, Harkness M, Knight R. Ratings of profile attractiveness after functional appliance treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 2000;**118**(4):371–6.
Webster T, Harkness M, Herbison P. Associations between changes in selected facial dimensions and the outcome of orthodontic treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 1996;**110**(1):46–53.
* Wijayaratne D, Harkness M, Herbison P. Functional appliance treatment assessed using the PAR index. *Australian Orthodontic Journal* 2000;**16**(3):118–26.
- North Carolina 2004** {published data only}
Almeida MA, Phillips C, Kula K, Tulloch C. Stability of the palatal rugae as landmarks for analysis of dental casts. *The Angle Orthodontist* 1995;**65**(1):43–8.
Brin I, Tulloch JF, Koroluk L, Phillips C. External apical root resorption in Class II malocclusion: a retrospective review of 1- versus 2-phase treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003;**124**(2): 151–6.
Dann C 4th, Phillips C, Broder HL, Tulloch JF. Self-concept, Class II malocclusion, and early treatment. *The Angle Orthodontist* 1995;**65**(6):411–6.
Ehmer U, Tulloch CJ, Proffit WR, Phillips C. An international comparison of early treatment of angle Class-II/1 cases. Skeletal effects of the first phase of a prospective clinical trial. *Journal of Orofacial Orthopedics* 1999;**60**(6): 392–408.
Koroluk LD, Tulloch JF, Phillips C. Incisor trauma and early treatment for Class II Division 1 malocclusion. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003; **123**(2):117–26.
Proffit WR, Tulloch JF. Preadolescent Class II problems: treat now or wait?. *American Journal of Orthodontics and Dentofacial Orthopedics* 2002;**121**(6):560–2.
Tulloch JF. Early versus late treatment for Class II malocclusions. *European Journal of Orthodontics* 1999;**21**(4):453.
Tulloch JF, Phillips C, Koch G, Proffit WR. The effect of early intervention on skeletal pattern in Class II malocclusion: a randomized clinical trial. *American Journal of Orthodontics and Dentofacial Orthopedics* 1997;**111**(4): 391–400.
Tulloch JF, Phillips C, Proffit WR. Benefit of early Class II treatment: progress report of a two-phase randomized clinical trial. *American Journal of Orthodontics and Dentofacial Orthopedics* 1998;**113**(1):62–72.
Tulloch JF, Proffit WR, Phillips C. Influences on the outcome of early treatment for Class II malocclusion. *American Journal of Orthodontics and Dentofacial Orthopedics* 1997;**111**(5):533–42.
* Tulloch JF, Proffit WR, Phillips C. Outcomes in a 2-phase randomized clinical trial of early Class II treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 2004;**125**(6):657–67.
Tulloch JF, Rogers L, Phillips C. Early results from a randomized clinical trial of growth modification in Class II malocclusion. *Journal of Dental Research* 1992;**72**(Spec Issue (IADR Abstracts)):523.
- Showkatbakhsh 2011** {published and unpublished data}
Showkatbakhsh R, Meybodi SE, Jamilian A, Meybodi SA, Meybodi EM. Treatment effects of R-appliance and anterior inclined bite plate in class II, division I malocclusion. *Journal of Applied Oral Science* 2011;**19**(6):634–8.
- Thiruvengkatachari 2010** {published data only}
Thiruvengkatachari B, Sandler J, Murray A, Walsh T, O'Brien K. Comparison of Twin-block and Dynamax appliances for the treatment of Class II malocclusion in adolescents: a randomized controlled trial. *American Journal of Orthodontics and Dentofacial Orthopedics* 2010; **138**(2):144.e1-9; discussion 144-5.
- UK (11-14) 2003** {published and unpublished data}
O'Brien K, Wright J, Conboy F, Sanjie Y, Mandall N, Chadwick S, et al. Effectiveness of treatment for Class II malocclusion with the Herbst or twin-block appliances: a randomized, controlled trial. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003;**124**(2): 128–37.
- UK (Mixed) 2009** {published and unpublished data}
O'Brien K, Wright J, Conboy F, Appelbe P, Davies L, Connolly I, et al. Early treatment for Class II Division 1 malocclusion with the Twin-block appliance: a multicenter, randomized, controlled trial. *American Journal of Orthodontics and Dentofacial Orthopedics* 2009;**135**(5): 573–9.
O'Brien K, Wright J, Conboy F, Chadwick S, Connolly I, Cook P, et al. Effectiveness of early orthodontic treatment with the Twin-block appliance: a multicenter, randomized, controlled trial. Part 2: Psychosocial effects. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003; **124**(5):488–94.
* O'Brien K, Wright J, Conboy F, Sanjie Y, Mandall N, Chadwick S, et al. Effectiveness of early orthodontic treatment with the Twin-block appliance: a multicenter,

randomized, controlled trial. Part 1: Dental and skeletal effects. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003;**124**(3):234–43.

Yaqoob 2012 {published data only}

Yaqoob O, Dibiasi AT, Cobourne MT. Use of the Clark twin block with and without a labial bow in the correction of class II division 1 malocclusion: a randomized clinical trial. *Journal of Orthodontics* 2010;**37**(4):310.

* Yaqoob O, Dibiasi AT, Fleming PS, Cobourne MT. Use of the Clark Twin Block functional appliance with and without an upper labial bow: a randomized controlled trial. *The Angle Orthodontist* 2012;**82**(2):363–9.

References to studies excluded from this review

Aknin 2000 {published data only}

Aknin JJ, Morra L. Comparative study of mandibular growth and rotation in two sample groups treated according to the “Distal Active Concept” or the Edgewise technique. *L'Orthodontie Francaise* 2000;**71**(4):343–61.

Ashmore 2002 {published data only}

Ashmore JL, Kurland BF, King GJ, Wheeler TT, Ghafari J, Ramsay DS. A 3-dimensional analysis of molar movement during headgear treatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 2002;**121**(1):18–30.

Baccetti 2009 {published data only}

Baccetti T, Franchi L, Kim LH. Effect of timing on the outcomes of 1-phase nonextraction therapy of Class II malocclusion. *American Journal of Orthodontics and Dentofacial Orthopedics* 2009;**136**(4):501–9.

Bishara 1995 {published data only}

Bishara SE, Cummins DM, Jakobsen JR, Zaher AR. Dentofacial and soft tissue changes in Class II, division 1 cases treated with and without extractions. *American Journal of Orthodontics and Dentofacial Orthopedics* 1995;**107**(1):28–37.

Chintakanon 2000 {published data only}

Chintakanon K, Sampson W, Wilkinson T, Townsend G. A prospective study of Twin-block appliance therapy assessed by magnetic resonance imaging. *American Journal of Orthodontics and Dentofacial Orthopedics* 2000;**118**(5):494–504.

Collett 2000 {published data only}

Collett AR. Current concepts on functional appliances and mandibular growth stimulation. *Australian Dental Journal* 2000;**45**(3):173–8.

Cura 1996 {published data only}

Cura N, Sarac M, Ozturk Y, Surmeli N. Orthodontic and orthopedic effects of Activator, Activator-HG combination, and Bass appliances: a comparative study. *American Journal of Orthodontics and Dentofacial Orthopedics* 1996;**110**(1):36–45.

Dahan 1989 {published data only}

Dahan J, Serhal JB, Englebert A. Cephalometric changes in Class II, Division 1 cases after orthopedic treatment with the bioactivator. *American Journal of Orthodontics and Dentofacial Orthopedics* 1989;**95**(2):127–37.

De Almeida 2002 {published data only}

De Almeida MR, Henriques JF, Ursi W. Comparative study of the Frankel (FR-2) and bionator appliances in the treatment of Class II malocclusion. *American Journal of Orthodontics and Dentofacial Orthopedics* 2002;**121**(5):458–66.

DeVincenzo 1989 {published data only}

DeVincenzo JP, Winn MW. Orthopedic and orthodontic effects resulting from the use of a functional appliance with different amounts of protrusive activation. *American Journal of Orthodontics and Dentofacial Orthopedics* 1989;**96**(3):181–90.

Du 2002 {published data only}

Du X, Hagg U, Rabie AB. Effects of headgear Herbst and mandibular step-by-step advancement versus conventional Herbst appliance and maximal jumping of the mandible. *European Journal of Orthodontics* 2002;**24**(2):167–74.

Erverdi 1995 {published data only}

Erverdi N, Ozkan G. A cephalometric investigation of the effects of the Elastic Bite-block in the treatment of Class II division 1 malocclusions. *European Journal of Orthodontics* 1995;**17**(5):375–84.

Falck 1989 {published data only}

Falck F, Frankel R. Clinical relevance of step-by-step mandibular advancement in the treatment of mandibular retrusion using the Frankel appliance. *American Journal of Orthodontics and Dentofacial Orthopedics* 1989;**96**(4):333–41.

Firouz 1992 {published data only}

Firouz M, Zernik J, Nanda R. Dental and orthopedic effects of high-pull headgear in treatment of Class II, division 1 malocclusion. *American Journal of Orthodontics and Dentofacial Orthopedics* 1992;**102**(3):197–205.

Franco 2002 {published data only}

Franco AA, Yamashita HK, Lederman HM, Cevidanes LH, Proffit WR, Vigorito JW. Frankel appliance therapy and the temporomandibular disc: a prospective magnetic resonance imaging study. *American Journal of Orthodontics and Dentofacial Orthopedics* 2002;**121**(5):447–57.

Freeman 2009 {published data only}

Freeman DC, McNamara JA Jr, Baccetti T, Franchi L, Frankel C. Long-term treatment effects of the FR-2 appliance of Frankel. *American Journal of Orthodontics and Dentofacial Orthopedics* 2009;**135**(5):570.e1–6; discussion 570–1.

Ghiglione 2000 {published data only}

Ghiglione V, Maspero C, Garagiola U. Skeletal Class II therapy - Effects of Bionator and Teuscher appliances. *European Journal Orthodontics* 2000;**22**(4):445.

Gianelly 1983 {published data only}

Gianelly AA, Brosnan P, Martignoni M, Bernstein L. Mandibular growth, condyle position and Frankel appliance therapy. *The Angle Orthodontist* 1983;**53**(2):131–42.

Guner 2003 {published data only}

Guner DD, Ozturk Y, Sayman HB. Evaluation of the effects of functional orthopaedic treatment on temporomandibular

- p>
joints with single-photon emission computerized tomography.
- European Journal of Orthodontics*
- 2003;
- 25**
- (1): 9–12.
- Hagg 2002** *{published data only}*
Hagg U, Tse EL, Rabie AB, Robinson W. A comparison of splinted and banded Herbst appliances: treatment changes and complications. *Australian Orthodontic Journal* 2002;**18**(2):76–81.
- Harvold 1971** *{published data only}*
Harvold EP, Vargervik K. Morphogenetic response to activator treatment. *American Journal of Orthodontics* 1971;**60**(5):478–90.
- Hiyama 2002** *{published data only}*
Hiyama S, Kuribayashi G, Ono T, Ishiwata Y, Kuroda T. Nocturnal masseter and suprahyoid muscle activity induced by wearing a bionator. *The Angle Orthodontist* 2002;**72**(1): 48–54.
- Ingervall 1991** *{published data only}*
Ingervall B, Thuer U. Temporal muscle activity during the first year of Class II, division 1 malocclusion treatment with an activator. *American Journal of Orthodontics and Dentofacial Orthopedics* 1991;**99**(4):361–8.
- Iscan 1997** *{published data only}*
Iscan HN, Sarisoy L. Comparison of the effects of passive posterior bite-blocks with different construction bites on the craniofacial and dentoalveolar structures. *American Journal of Orthodontics and Dentofacial Orthopedics* 1997;**112**(2):171–8.
- Janson 2003** *{published data only}*
Janson GR, Toruno JL, Martins DR, Henriques JF, de Freitas MR. Class II treatment effects of the Frankel appliance. *European Journal of Orthodontics* 2003;**25**(3): 301–9.
- Jarrell 2001** *{published data only}*
Jarrell KT, Hudson JM, Killiany DM. Activator-headgear combination appliance treatment of Class II, division I malocclusion. *Journal of Dental Research* 2001;**80** Special Issue:180 (Abs No 1156).
- Kalra 1989** *{published data only}*
Kalra V, Burstone CJ, Nanda R. Effects of a fixed magnetic appliance on the dentofacial complex. *American Journal of Orthodontics and Dentofacial Orthopedics* 1989;**95**(6): 467–78.
- Keski-Nisula 2003** *{published data only}*
Keski-Nisula K, Lehto R, Lusa V, Keski-Nisula L, Varrelä J. Occurrence of malocclusion and need of orthodontic treatment in early mixed dentition. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003;**124**(6): 631–8.
- Kumar 1996** *{published data only}*
Kumar S, Sidhu SS, Kharbanda OP. A cephalometric evaluation of the dental and facial-skeletal effects using the Bionator with stepwise protrusive activations. *The Journal of Clinical Pediatric Dentistry* 1996;**20**(2):101–8.
- Lange 1995** *{published data only}*
Lange DW, Kalra V, Broadbent BH Jr, Powers M, Nelson S. Changes in soft tissue profile following treatment with the bionator. *The Angle Orthodontist* 1995;**65**(6):423–30.
- Lund 1998** *{published data only}*
Lund DI, Sandler PJ. The effects of Twin Blocks: a prospective controlled study. *American Journal of Orthodontics and Dentofacial Orthopedics* 1998;**113**(1): 104–10.
- Malmgren 1987** *{published data only}*
Malmgren O, Omblus J, Hagg U, Pancherz H. Treatment with an orthopedic appliance system in relation to treatment intensity and growth periods. A study of initial effects. *American Journal of Orthodontics and Dentofacial Orthopedics* 1987;**91**(2):143–51.
- Malta 2010** *{published data only}*
Malta LA, Baccetti T, Franchi L, Faltin K Jr, McNamara JA Jr. Long-term dentoskeletal effects and facial profile changes induced by bionator therapy. *The Angle Orthodontist* 2010;**80**(1):10–7.
- Meral 2004** *{published data only}*
Meral O, Iscan HN, Okay C, Gursoy Y. Effects of bilateral upper first premolar extraction on the mandible. *European Journal of Orthodontics* 2004;**26**(2):223–31.
- Muniandy 2000** *{published data only}*
Muniandy SD, Battagel JM, Moss JP. A prospective study of the twin block and silensor appliances. *European Journal of Orthodontics* 2000;**22**(5):604.
- Nelson 2000** *{published data only}*
Nelson B, Hansen K, Hagg U. Class II correction in patients treated with class II elastics and with fixed functional appliances: a comparative study. *American Journal of Orthodontics and Dentofacial Orthopedics* 2000;**118**(2): 142–9.
- Op Heij 1989** *{published and unpublished data}*
Op Heij DG, Callaert H, Opdebeeck HM. The effect of the amount of protrusion built into the bionator on condylar growth and displacement: a clinical study. *American Journal of Orthodontics and Dentofacial Orthopedics* 1989;**95**(5): 401–9.
- Ozturk 1994** *{published data only}*
Ozturk Y, Tankuter N. Class II: a comparison of activator and activator headgear combination appliances. *European Journal of Orthodontics* 1994;**16**(2):149–57.
- Pangrazio 1999** *{published data only}*
Pangrazio-Kulbersh V, Kaczynski R, Shunock M. Early treatment outcome assessed by the Peer Assessment Rating index. *American Journal of Orthodontics and Dentofacial Orthopedics* 1999;**115**(5):544–50.
- Pangrazio 2003** *{published data only}*
Pangrazio-Kulbersh V, Berger JL, Chermak DS, Kaczynski R, Simon ES, Haerian A. Treatment effects of the mandibular anterior repositioning appliance on patients with Class II malocclusion. *American Journal of Orthodontics and Dentofacial Orthopedics* 2003;**123**(3):286–95.

Parkin 2001 {published data only}

Parkin NA, McKeown HF, Sandler PJ. Comparison of 2 modifications of the twin-block appliance in matched Class II samples. *American Journal of Orthodontics and Dentofacial Orthopedics* 2001;**119**(6):572–7.

Phan 2006 {published data only}

Phan KL, Bendeus M, Hagg U, Hansen K, Rabie AB. Comparison of the headgear activator and Herbst appliance-effects and post-treatment changes. *European Journal of Orthodontics* 2006;**28**(6):594–604.

Pirttiniemi 2005 {published data only}

Mantysaari R, Kantomaa T, Pirttiniemi P, Pykalainen A. The effects of early headgear treatment on dental arches and craniofacial morphology: a report of a 2 year randomized study. *European Journal of Orthodontics* 2004;**26**(1):59–64.
* Pirttiniemi P, Kantomaa T, Mantysaari R, Pykalainen A. The effects of early headgear treatment on dental arches and craniofacial morphology: an 8 year report of a randomized study. *European Journal of Orthodontics* 2005;**27**(5):429–36.

Reukers 1998 {published data only}

Reukers EA, Sanderink GC, Kuijpers-Jagtman AM, van't Hof MA. Radiographic evaluation of apical root resorption with 2 different types of edgewise appliances. Results of a randomized clinical trial. *Journal of Orofacial Orthopedics* 1998;**59**(2):100–9.

Sari 2003 {published data only}

Sari Z, Goyenc Y, Doruk C, Usumez S. Comparative evaluation of a new removable Jasper Jumper functional appliance vs an activator-headgear combination. *The Angle Orthodontist* 2003;**73**(3):286–93.

Schaefer 2004 {published data only}

Schaefer AT, McNamara JA Jr, Franchi L, Baccetti T. A cephalometric comparison of treatment with the Twin-block and stainless steel crown Herbst appliances followed by fixed appliance therapy. *American Journal of Orthodontics and Dentofacial Orthopedics* 2004;**126**(1):7–15.

Shannon 2004 {published data only}

Shannon KR, Nanda RS. Changes in the curve of Spee with treatment and at 2 years posttreatment. *American Journal of Orthodontics and Dentofacial Orthopedics* 2004;**125**(5):589–96.

Siara-Olds 2010 {published data only}

Siara-Olds NJ, Pangrazio-Kulbersh V, Berger J, Bayirli B. Long-term dentoskeletal changes with the Bionator, Herbst, Twin Block, and MARA functional appliances. *The Angle Orthodontist* 2010;**80**(1):18–29.

Siqueira 2007 {published data only}

Siqueira DF, de Almeida RR, Janson G, Brandao AG, Coelho Filho CM. Dentoskeletal and soft-tissue changes with cervical headgear and mandibular protraction appliance therapy in the treatment of Class II malocclusions. *American Journal of Orthodontics and Dentofacial Orthopedics* 2007;**131**(4):447.e21–30.

Taner 2003 {published data only}

Taner TU, Yukay F, Pehlivanoglu M, Cakirer B. A comparative analysis of maxillary tooth movement produced by cervical headgear and pend-x appliance. *The Angle Orthodontist* 2003;**73**(6):686–91.

Thuer 1989 {published data only}

Thuer U, Ingervall B, Burgin W. Does the mandible alter its functional position during activator treatment?. *American Journal of Orthodontics and Dentofacial Orthopedics* 1989;**96**(6):477–84.

Tumer 1999 {published data only}

Tumer N, Gultan AS. Comparison of the effects of monoblock and twin-block appliances on the skeletal and dentoalveolar structures. *American Journal of Orthodontics and Dentofacial Orthopedics* 1999;**116**(4):460–8.

Ucem 1998 {published data only}

Ucem TT, Yuksel S. Effects of different vectors of forces applied by combined headgear. *American Journal of Orthodontics and Dentofacial Orthopedics* 1998;**113**(3):316–23.

Ucuncu 2001 {published and unpublished data}

Ucuncu N, Turk T, Carels C. Comparison of modified Teuscher and van Beek functional appliance therapies in high-angle cases. *Journal of Orofacial Orthopedics* 2001;**62**(3):224–37.

Wey 2007 {published data only}

Wey MC, Bendeus M, Peng L, Hagg U, Rabie AB, Robinson W. Stepwise advancement versus maximum jumping with headgear activator. *European Journal of Orthodontics* 2007;**29**(3):283–93.

Wieslander 1984 {published data only}

Wieslander L. Intensive treatment of severe Class II malocclusions with a headgear-Herbst appliance in the early mixed dentition. *American Journal of Orthodontics* 1984;**86**(1):1–13.

Witt 1999 {published data only}

Witt E, Watted N. Effectiveness of intra- and extraoral aids to the bionator. A controlled study within the scope of the “Wuerzburg concept”. *Journal of Orofacial Orthopedics* 1999;**60**(4):269–78.

Additional references

Antonarakis 2007

Antonarakis GS, Kiliaridis S. Short-term anteroposterior treatment effects of functional appliances and extraoral traction on class II malocclusion. A meta-analysis. *The Angle Orthodontist* 2007;**77**(5):907–14.

Barnett 2008

Barnett GA, Higgins DW, Major PW, Flores-Mir C. Immediate skeletal and dentoalveolar effects of the crown- or banded type Herbst appliance on Class II division 1 malocclusion. *The Angle Orthodontist* 2008;**78**(2):361–9.

Cozza 2006

Cozza P, Baccetti T, Franchi L, De Toffol L, McNamara JA Jr. Mandibular changes produced by functional appliances

- in Class II malocclusion: a systematic review. *American Journal of Orthodontics and Dentofacial Orthopedics* 2006;**129**(5):599.e1-12; discussion e1-6.
- El-Mangoury 1990**
El-Mangoury NH, Mostafa YA. Epidemiologic panorama of dental occlusion. *The Angle Orthodontist* 1990;**60**(3): 207–14.
- Flores-Mir 2007**
Flores-Mir C, Aye A, Goswami A, Charkhandeh S. Skeletal and dental changes in Class II division 1 malocclusions treated with splint-type Herbst appliances. A systematic review. *The Angle Orthodontist* 2007;**77**(2):376–81.
- Higgins 2011**
Higgins JPT, Green S (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available from www.cochrane-handbook.org.
- Holmes 1992**
Holmes A. The prevalence of orthodontic treatment need. *British Journal of Orthodontics* 1992;**19**(3):177–82.
- Jahal 2007**
Jahal A, Cheung MY, Marcene W. The impact of two different malocclusion traits on quality of life. *British Dental Journal* 2007;**202**:E2.
- Landis 1977**
Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;**33**(1): 159–74.
- Nguyen 1999**
Nguyen QV, Bezemer PD, Habets L, Prahl-Andersen B. A systematic review of the relationship between overjet size and traumatic dental injuries. *European Journal of Orthodontics* 1999;**21**(5):503–15.
- Perillo 2012**
Perillo L, Cannavale R, Ferro F, Franchi L, Masucci C, Chiodini P, et al. Meta-analysis of skeletal mandibular changes during Frankel appliance treatment. *European Journal of Orthodontics* 2012;**33**(1):84–92.
- Proffit 1993**
Proffit WR, Fields HW Jr. *Contemporary Orthodontics*. 2nd Edition. St Louis, USA: Mosby-Year Book, Inc, 1993:2–16.
- Seehra 2011**
Seehra J, Fleming PS, Newton T, DiBiase AT. Bullying in orthodontic patients and its relationship to malocclusion, self-esteem and oral health-related quality of life. *Journal of Orthodontics* 2011;**38**(4):247–56.
- Shaw 1980**
Shaw WC, Addy M, Ray C. Dental and social effects of malocclusion and effectiveness of orthodontic treatment: a review. *Community Dentistry and Oral Epidemiology* 1980;**8** (1):36–45.
- Shaw 1991**
Shaw WC, Richmond S, O'Brien KD, Brook P, Stephens CD. Quality control in orthodontics: indices of treatment need and treatment standards. *British Dental Journal* 1991;**170**(3):107–12.
- Silva 2001**
Silva RG, Kang DS. Prevalence of malocclusion among Latino adolescents. *American Journal of Orthodontics and Dentofacial Orthopedics* 2001;**119**(3):313–5.
- * Indicates the major publication for the study